Rencontres de Moriond EW 2014



ID de Contribution: 191

Type: YSF (Young Scientists Forum)

High SUSY scale and a few-parameter description of the scalar sector

vendredi 21 mars 2014 19:38 (5 minutes)

We analyze the Minimal Supersymmetric extension of the Standard Model that we have after the discovery of the Higgs boson at the LHC, i.e. a model in which the lighter h boson has a mass of approximately 125 GeV which, together with the non-observation of superparticles at the LHC, indicates that the SUSY-breaking scale MS is rather high. We first demonstrate that the value Mh≈125 GeV fixes the dominant radiative corrections that enter the MSSM Higgs boson masses, leading to a Higgs sector that can be described, to a good approximation, by only two free parameters. In a second step, we consider the direct supersymmetric radiative corrections and show that, to a good approximation, the phenomenology of the lighter Higgs state can be described by its mass and three couplings : those to massive gauge bosons and to top and bottom quarks. We perform a fit of these couplings using the latest LHC data on the production and decay rates of the light h boson and combine it with the limits from the negative search of the heavier H,A and H± states.

Auteur principal:QUEVILLON, Jérémie (LPT)Orateur:QUEVILLON, Jérémie (LPT)Classification de Session:Young Scientist Forum 4

Classification de thématique: Theory