## **Rencontres de Moriond EW 2014**



ID de Contribution: 154

Type: Ordinary

## Enhanced diphoton signal of a light singlet-like scalar in NMSSM

vendredi 21 mars 2014 11:15 (15 minutes)

NMSSM with a light singlet-like scalar and strongly suppressed couplings to b and  $\tau$  is investigated. It is shown that in such a scenario the singlet-like scalar to diphoton signal can be larger than for the SM Higgs for a wide range of masses between 60 and 110 GeV, in agreement with all the LEP and LHC data. Enhancement of the singlet-like scalar to diphoton signal is correlated with positive correction to the SM-like Higgs mass from mixing between SM-like Higgs and the singlet. It is also shown that the couplings to b and  $\tau$  and, in consequence, branching ratios of the SM-like Higgs are anti-correlated with those of the singlet-like scalar. If the singlet-like scalar to diphoton signal is enhanced, the signal strengths of the 125 GeV Higgs in the diphoton and WW\*/ZZ\* channels are predicted to be smaller than for the SM Higgs.

**Auteurs principaux:** M. BADZIAK, Marcin (University of Warsaw); OLECHOWSKI, Marek (University of Warsaw); POKORSKI, Stefan (University of Warsaw)

**Orateur:** M. BADZIAK, Marcin (University of Warsaw)

Classification de Session: Beyond the Standard Model

Classification de thématique: Theory