



Oliver Buchmueller, Imperial College London

SEARCHES FOR DARK MATTER PRODUCTION AT THE LHC





Fundamental Open Questions in Particle Physics

SUSY & DM Searches, @ LHC O. Buchmüller



I. What is the origin of mass?

- Why are the vector bosons Z and W massive whereas the photon is massless?
- Is there a Higgs boson - or even more of them ?





I. Is there a new symmetry - Supersymmetry ?

- Can we get experimental evidence to support the Grand Unification of all fundamental forces?
- What is the origin of Dark Matter in the Universe?
→ Is a fundamental particle responsible for it?



III. What is the origin of the matter-anti-matter asymmetry in our Universe?



- Does the answer lie in CP violation?
- Neutrino masses and mixing - how do they fit in the picture?



Fundamental Open Questions in Particle Physics



I. What is the origin of mass?

- Why are the vector bosons Z and W are massive whereas the photon is massless?
- Is there a Higgs boson - or even more of them ?





I. Is there a new symmetry - Supersymmetry ?

- Can we get experimental evidence to support the Grand Unification of all fundamental forces?
- What is the origin of Dark Matter in the Universe?
→ Is a fundamental particle responsible for it?



III. What is the origin of the matter-anti-matter asymmetry in our Universe?

- Does the answer lie in CP violation?
- Neutrino masses and mixing - how do they fit in the picture?




The Large Hadron Collider at CERN

SUSY & DM Searches, @ LHC O. Buchmüller




The Large Hadron Collider at CERN

SUSY & DM Searches, @ LHC O. Buchmüller





The Large Hadron Collider at CERN

LHC : 27 km long
100m underground




General Purpose,
pp, heavy ions




The Large Hadron Collider at CERN


LHC : 27 km long
100m underground




General Purpose,
pp, heavy ions



Heavy ions, pp



The Large Hadron Collider at CERN



Searches for SUSY (& DM)




Searches for SUSY (& DM)




Supersymmetry

Extension of the Standard Model: Introduce a new symmetry
Spin 1/2 matter particles (fermions) $\Leftrightarrow$ Spin 1 force carriers (bosons)

Standard Model particles



SUSY particles




New Quantum number: R-parity: $R_p = (-1)^{B+L+2s} = +1$ SM particles
R-parity conservation: -1 SUSY particles

- SUSY particles are produced in pairs
- The lightest SUSY particle (LSP) is stable

What do we call a “SUSY search”?

*The definition is purely derived from the experimental signature.
Therefore, a “SUSY search signature” is characterized by*

Lots of missing energy, many jets, and possibly leptons in the final state



Missing Energy:

- from LSP

Multi-Jet:

- from cascade decay (gaugino)

Multi-Leptons:


- from decay of charginos/neutralinos

RP-Conserving SUSY is a very prominent example predicting this famous signature but ...

What is its experimental signature?

SUSY & DM Searches, @ LHC O. Buchmüller

... by no means is it the only New Physics model predicting this experimental pattern. Many other NP models predict this genuine signature



Missing Energy:

- *Nwimp* - end of the cascade

Multi-Jet:


- from decay of the *N*s (possibly via heavy SM particles like top, W/Z)

Multi-Leptons:

- from decay of the *N*'s

Model examples are Extra dimensions, Little Higgs, Technicolour, etc
but a more generic definition for this signature is as follows.

Early SUSY Search Strategy at the LHC



Search Signatures


- SUSY-like decay chains range from short to long and simple to very complicated.
- All physics objects, MET, jets, leptons, photons, b's taus, tops, W, Z, etc are involved
- Comprehensive coverage of all possible signature requires a topology oriented search strategy:

References Analyses

0-leptons	1-lepton	OSDL	SSDL	≥ 3 leptons	2-photons	γ +lepton
Jets + MET	Single lepton + Jets + MET	Opposite-sign di-lepton + jets + MET	Same-sign di-lepton + jets + MET	Multi-lepton	Di-photon + jet + MET	Photon + lepton + MET


Already in less then two years of operation ATLAS & CMS managed to carry out the full list of these core “SUSY References Analyses”!

Inclusive SUSY Searches in 2013




Inclusive SUSY Searches in 2013

SUSY & DM Searches, @ LHC O. Buchmüller



Inclusive SUSY Searches in 2013


SUSY & DM Searches, @ LHC O. Buchmüller



The LHC has pushed the mass scale in constraint SUSY models to a new level!

CMSSM: Evolution with time

SUSY & DM Searches, @ LHC O. Buchmüller



χ^2 increase from
bluish to reddish



Source:

<http://mastercode.web.cern.ch/mastercode/>

Observable	Source Th./Ex.	Constraint	$\Delta\chi^2$ (CMSSM)	$\Delta\chi^2$ (NUHM1)	$\Delta\chi^2$ ("SM")
m_t [GeV]	[43]	173.2 ± 0.90	0.05	0.06	-
$\Delta\alpha_{had}^{(5)}$ (M_Z)	[42]	0.02749 ± 0.00010	0.009	0.004	-
M_Z [GeV]	[44]	91.1875 ± 0.0021	2.7×10^{-10}	0.26	-
Γ_Z [GeV]	[26] / [44]	$2.4952 \pm 0.0023 \pm 0.001_{SUSY}$	0.078	0.047	0.14
σ_{had} [nb]	[26] / [44]	41.540 ± 0.037	2.50	2.57	2.54
R_l	[26] / [44]	20.767 ± 0.025	1.05	1.08	1.08
$A_{fb}(\ell)$	[26] / [44]	0.01714 ± 0.00095	0.72	0.69	0.81
$A_T(P_T)$	[26] / [44]	0.1465 ± 0.0032	0.11	0.13	0.07
R_b	[26] / [44]	0.21629 ± 0.00066	0.26	0.29	0.27
R_c	[26] / [44]	0.1721 ± 0.0030	0.002	0.002	0.002
$A_{fb}(b)$	[26] / [44]	0.0992 ± 0.0016	7.17	7.37	6.63
$A_{fb}(c)$	[26] / [44]	0.0707 ± 0.0035	0.86	0.88	0.80
A_b	[26] / [44]	0.923 ± 0.020	0.36	0.36	0.35
A_c	[26] / [44]	0.670 ± 0.027	0.005	0.005	0.005
A_ℓ (SLD)	[26] / [44]	0.1513 ± 0.0021	3.16	3.03	3.51
$\sin^2 \theta_C^{\ell}(Q_{FB})$	[26] / [44]	0.2324 ± 0.0012	0.63	0.64	0.59
M_W [GeV]	[26] / [44]	$80.399 \pm 0.023 \pm 0.010_{SUSY}$	1.77	1.39	2.08
$a_\mu^{had} - a_\mu^{SM}$	[53] / [42,54]	$(30.2 \pm 8.8 \pm 2.0_{SUSY}) \times 10^{-10}$	4.35	1.82	11.19 (N/A)
M_h [GeV]	[28] / [55,56]	$> 114.4[\pm 1.5_{SUSY}]$	0.0	0.0	0.0
$BR_{b \rightarrow sy}^{EXPT/SM}$	[45] / [46]	$1.117 \pm 0.076_{EXP} \pm 0.082_{SM} \pm 0.050_{SUSY}$	1.83	1.09	0.94
$BR(B_s \rightarrow \mu^+ \mu^-)$	[29] / [41]	CMS & LHCb	0.04	0.44	0.01
$BR_{B \rightarrow \tau \nu}^{EXPT/SM}$	[29] / [46]	$1.43 \pm 0.43_{EXP+TH}$	1.43	1.59	1.00
$BR(B_d \rightarrow \mu^+ \mu^-)$	[29] / [46]	$< 4.6[\pm 0.01_{SUSY}] \times 10^{-9}$	0.0	0.0	0.0
$BR_{K \rightarrow \mu \nu}^{EXPT/SM}$	[47] / [46]	0.99 ± 0.32	0.02	$\ll 0.01$	$\ll 0.01$
$BR_{K \rightarrow \pi \nu}^{EXPT/SM}$	[29] / [48]	$1.008 \pm 0.014_{EXP+TH}$	0.39	0.42	0.33
$BR_{K \rightarrow \pi \nu}^{EXPT/SM}$	[49] / [50]	< 4.5	0.0	0.0	0.0
$\Delta M_{B_s}^{EXPT/SM}$	[49] / [51,52]	$0.97 \pm 0.01_{EXP} \pm 0.27_{SM}$	0.02	0.02	0.01
$\Delta M_{B_d}^{EXPT/SM}$	[29] / [46,51,52]	$1.00 \pm 0.01_{EXP} \pm 0.13_{SM}$	$\ll 0.01$	0.33	$\ll 0.01$
$\Delta M_{K_S}^{EXPT/SM}$	[49] / [51,52]	$1.08 \pm 0.14_{EXP+TH}$	0.27	0.37	0.33
$\Omega_{CDM} h^2$	[31] / [13]	$0.1120 \pm 0.0056 \pm 0.012_{SUSY}$	8.4×10^{-4}	0.1	N/A
$\sigma_p^{2\gamma}$	[25]	$(m_{\chi_{1,2}^0}, \sigma_p^{2\gamma})$ plane	0.13	0.13	N/A
jets + B_T	[18,20]	$(m_0, m_{1/2})$ plane	1.55	2.20	N/A
$H/A, H^\pm$	[21]	$(M_A, \tan \beta)$ plane	0.0	0.0	N/A
Total $\chi^2/d.o.f.$	All	All	28.8/22	27.3/21	32.7/23 (21.5/22)
p-values			15%	16%	9% (49%)

Global Fit to indirect and direct constraints on SUSY!





Other “fitter” groups find very similar results: e.g.

SuperBayeS: [arXiv:1212.2636](https://arxiv.org/abs/1212.2636)

Fittino group: [arXiv:1204.4199](https://arxiv.org/abs/1204.4199)

CMSSM: Evolution with time


SUSY & DM Searches, @ LHC O. Buchmüller



SUSY Status – post 7 TeV LHC data

- Constrained SUSY models like the CMSSM are severely put under pressure by the LHC limits!
- Experiments define new benchmarks and less complex SUSY models in order to present the interpretation of their searches.
- Aided by the discovery of a Higgs boson, the focus of the experimental search strategy and corresponding interpretation shifts towards other scenarios like “Natural SUSY” (i.e. 3rd generation squark searches).


What the individual searches are sensitive to is much more simple...



SMS: a few interesting features


SUSY & DM Searches, @ LHC O. Buchmüller

$m_{\text{LSP}}^{\text{max}} \approx 0.3 \text{ TeV}$: LSP mass above
which there is NO limit anymore



$m_G^{\text{max}} \approx 0.8 \text{ TeV}$: Best limit in plane

Assumes 100%
BR for decay chain
considered.



$$\tilde{q}\tilde{q} \rightarrow q\tilde{\chi}^0\bar{q}\tilde{\chi}^0$$


How to summarize SMS limits?

*Approach taken in the 2012 and 2013 Experimental SUSY PDG reviews
[OB & Paul De Jong]:*

<http://pdg.lbl.gov/2012/reviews/rpp2012-rev-susy-2-experiment.pdf>

<http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf>

Model	Assumption	$m_{\tilde{q}}$	$m_{\tilde{g}}$
CMSSM	$m_{\tilde{q}} \approx m_{\tilde{g}}$	1400	1400
	all $m_{\tilde{q}}$	-	800
	all $m_{\tilde{g}}$	1300	-
Simplified model $\tilde{g}\tilde{g}$	$m_{\tilde{\chi}_1^0} = 0$	-	900
	$m_{\tilde{\chi}_1^0} > 300$	-	no limit
Simplified model $\tilde{q}\tilde{q}$	$m_{\tilde{\chi}_1^0} = 0$	750	-
	$m_{\tilde{\chi}_1^0} > 250$	no limit	-
Simplified model $\tilde{g}\tilde{q}, \tilde{g}\tilde{\bar{q}}$	$m_{\tilde{\chi}_1^0} = 0, m_{\tilde{q}} \approx m_{\tilde{g}}$	1500	1500
	$m_{\tilde{\chi}_1^0} = 0, \text{ all } m_{\tilde{g}}$	1400	-
	$m_{\tilde{\chi}_1^0} = 0, \text{ all } m_{\tilde{q}}$	-	900



This was an appropriate approach for the rather limited amount of inclusive searches and corresponding SMS interpretations available in 2011 (7 TeV).

How to summarize SMS limits?

*Approach taken in the 2012 and 2013 Experimental SUSY PDG reviews
[OB & Paul De Jong]:*


<http://pdg.lbl.gov/2012/reviews/rpp2012-rev-susy-2-experiment.pdf>

<http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf>


Model	Assumption	$m_{\tilde{q}}$	$m_{\tilde{g}}$
	$m_{\tilde{q}} \approx m_{\tilde{g}}$	1400	1400
CMSSM	all m	800	800

It is a challenge to do justice to the many searches and limits that have been established so far
- even more so to put it all together into the/a "bigger picture".

Simplified model $\tilde{q}\tilde{q}$	$m_{\tilde{\chi}_1^0} = 0$	750	-
	$m_{\tilde{\chi}_1^0} > 250$	no limit	-
Simplified model $\tilde{g}\tilde{q}, \tilde{g}\tilde{\tilde{q}}$	$m_{\tilde{\chi}_1^0} = 0, m_{\tilde{q}} \approx m_{\tilde{g}}$	1500	1500
	$m_{\tilde{\chi}_1^0} = 0, \text{all } m_{\tilde{g}}$	1400	-
	$m_{\tilde{\chi}_1^0} = 0, \text{all } m_{\tilde{q}}$	-	900



This was an appropriate approach for the rather limited amount of inclusive searches and corresponding SMS interpretations available in 2011 (7 TeV).



m_{LSP}
[GeV]

1000

750

500

250

0

0

250

500

750

1000

1250

1500


26

Note: The following results are based
on PDG review September 2013.

<http://pdg.lbl.gov/2013/reviews/rpp2013-rev-susy-2-experiment.pdf>

$m_{\text{SUSY}} - m_{\text{LSP}} < m_t$


What are representative
SMS limits on the different
particles?



m_{SUSY}
[GeV]

Direct squark production – chosen limits

SUSY & DM Searches, @ LHC O. Buchmüller



CMS-SUS-PAS-13-012


Signature: Jets + E_T^{miss} + H_T
Limit assumes all 1st & 2nd gen squarks to be mass degenerate or only one light squark!


ATLAS-CONF-2013-053


Signature: 2 b-jets + E_T^{mis}

ATLAS-CONF-2013-037

Signature: 1Lepton + jets + E_T^{mis}









Gluino mediated squark production – limits chosen



© LHC O. Buchmüller



CMS Preliminary, 19.5 fb⁻¹, $\sqrt{s} = 8$ TeV




CMS, L = 19.4 fb⁻¹, $\sqrt{s} = 8$ TeV




CMS-SUS-PAS-13-012

Signature: Jets + H_T + E_t^{miss}




CMS-SUS-PAS-12-024

Signature: : Jets + b-tag + E_t^{miss}



ATLAS-CONF-2013-061

Signature: 0/1 Leptons +
3 b-tag + E_t^{mis}



Gluino mediated:

Similar to direct squark production;
better control of background via
b-tagging and special topology
signatures (e.g. many jets) provide
higher sensitivity on gluino decay
chains involving stops and sbottoms.

Gluino mediated

$$m_{SUSY} = m_{\tilde{g}}$$

$$\tilde{g} \rightarrow q\bar{q}\chi_1^0$$


CMS-PAS-SUS-13-012


$$\tilde{g} \rightarrow b\bar{b}\chi_1^0$$

CMS-PAS-SUS-12-024

$$\tilde{g} \rightarrow t\bar{t}\chi_1^0$$



ATLAS-CONF-2013-061





Dedicated searches for direct stop-pair production



SUSY & DM Searches, @ LHC O. Buchmüller



LEP:
LEPSUSY
WG/04-01.1


CDF:
1203.4171
D0:
0803.2263


Status: July 2013





Dedicated searches for direct stop-pair production

SUSY & DM Searches, @ LHC O. Buchmüller















Direct slepton production




ATLAS-CONF-2013-049
 Signature
 2 lepton + E_T^{miss}







Direct chargino/neutralino production




light
slepton
"easy"


CMS-PAS-SUS-13-006
3l OSSF categorised in
bins of M_{ll} , MT, and ETmis









What does this imply for Linear Collider?






Dark Matter Searches: Direct Detection Experiments

SUSY & DM Searches, @ LHC O. Buchmüller



Dark Matter Searches: Direct Detection Experiments


SUSY & DM Searches, @ LHC O. Buchmüller



Also the low
Wimp mass region
($< \sim 20$ GeV)
is very interesting!


Direct Detection Landscape in a nutshell!

SUSY & DM Searches, @ LHC O. Buchmüller




Direct Detection Landscape in a nutshell!

SUSY & DM Searches, @ LHC O. Buchmüller




SUSY & Dark Matter: Evolution with time

SUSY & DM Searches, @ LHC O. Buchmüller




SUSY & Dark Matter: Evolution with time

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter Searches

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter Searches

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter Searches

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter Searches

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter Searches: Direct Detection vs Colliders

SUSY & DM Searches, @ LHC O. Buchmüller




Direct Detection Experiments

- DM-nucleus scattering




Collider Experiments

- Pair-production of DM
- missing energy signature




Mono-Mania (at the LHC)


Mono-Z




Mono-photon




Mono-jet




Mono-Higgs



Mono-W




Mono-top




Mono-Mania (at the LHC)


Mono-Z




Mono-photon




Mono-jet




Mono-Higgs



Mono-W





Mono-top



Monojet analyses better than direct detection?!

SUSY & DM Searches, @ LHC O. Buchmüller



Claim [often made]:


For **low mass** and the entire **spin-dependent** case monojet limits are stronger than direct detection limits!

Effective Field Theory (EFT) Interpretation

Example of considered operators:

$$O_V = \frac{(\bar{\chi}\gamma_\mu\chi)(\bar{q}\gamma_\mu q)}{\Lambda^2} \quad \text{Vector operator, s-channel}$$

$$O_{AV} = \frac{(\bar{\chi}\gamma_\mu\gamma_5\chi)(\bar{q}\gamma_\mu\gamma_5 q)}{\Lambda^2} \quad \text{Axial vector operator, s-channel}$$



Assumption of EFT


If the operator (e.g. V or AV) mediator is **suitably(!!)** heavy it can be integrated out to obtain the effective V or AV contact operator. **In this case (and only this case)**, the contact interaction scale Λ is related to the parameters entering the Lagrangian:

$$\Lambda = \frac{M_{mediator}}{\sqrt{g_q g_\chi}} \quad (\text{relation in the full theory})$$


Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan, C.McCabe: arXiv:1308.6799

➤ Compare Effective Field Theory (EFT) with Full Theory (FT)



Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Compare prediction of FT with EFT in $m_{\text{med}} - m_{\text{DM}}$ plane. Three regions become visible:

Region I: EFT and FT agree better than 20%

➤ EFT is valid!

Region II: EFT yields significant weaker limits than FT

➤ EFT limits are too conservative!


Region III: EFT yields significant stronger limits than FT

➤ EFT limits are too aggressive!


Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan, C.McCabe: arXiv:1308.6799

➤ Compare Effective Field Theory (EFT) with Full Theory (FT)



Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Three Regions as function of mediator mass:

Region I: Heavy m_{med}

➤ EFT is valid!

Region II: Medium m_{med} – Resonant enhancement

➤ EFT limits are too conservative!


Region III: Low m_{med}

➤ EFT limits are too aggressive!

Minimal Simplified Dark Matter Model

SUSY & DM Searches, @ LHC O. Buchmüller

Based on work from :
OB, S. Malik,
M.Dolan,C.McCabe



s-channel

Define simplified model with
(minimum) 4 parameters

Mediator mass (M_{med})	DM mass (M_{DM})
g_q	g_{DM}

DM

Dirac fermion	Scalar - real
Majorana fermion	Scalar - complex



Consider comprehensive set
of diagrams for mediator

Vector	Axial-vector
Scalar	Pseudoscalar

Dark Matter Limits from Direct Searches: Today

Examples: CMS monojet search
and recent LUX result:
interpretation in simplified models

SUSY & DM Searches, @ LHC O. Buchmüller




Assumes:

- couplings to all quarks
- $g_q = g_{DM} = 1$
- width calculated from g


**Direct Detection experiments
and collider are complementary
probing different regions of
the relevant parameter space!**

The Vector Case




The Vector Case

SUSY & DM Searches, @ LHC O. Buchmüller




While for Axial-Vector collider and DD are complementary in the full parameter space, for the pure Vector case, DD limits are strong EXCEPT for low DM masses!




Note: Vector scales like Atomic number squared ($\sim 130^2$) while AV scales like the spin (~ 1).

Future Projections



Compare:
LHC @ 300/fb and HL-LHC@/3000/fb
with Direct Detection:
Lux-Zeppline (~2025) and Neutrino noise border

Future Projections



Important complementarity of the two experimental approaches will allow good coverage of the relevant parameter space!

Big discovery potential!

Outlook: 8 TeV vs 14 TeV

SUSY & DM Searches, @ LHC O. Buchmüller

Use parton luminosities to illustrate the gain of 14 vs 8 TeV

Higgs:

$pp \rightarrow H$, $H \rightarrow WW$, ZZ and $\gamma\gamma$
mainly gg : factor ~ 2

SUSY – 3rd Generation:

Mass scale ~ 500 GeV

qq and gg : factor ~ 3 to 6

SUSY – Squarks/Gluino:


Mass scale ~ 1.5 TeV

qq, gg, qg : factor ~ 40 to 80

Z' :

Mass scale ~ 5 TeV

qq : factor ~ 1000




Increase in energy will help a lot!
Not just for SUSY...


Summary


- So far New Physics has not revealed itself!
 - Even by 2010 the LHC has entered new territory for New Physics searches and since pushed e.g. the (coloured) SUSY mass scale to the ~ 1 TeV scale
 - We were well prepared for an early discovery but we also knew that it could take more time and ingenuity before we can claim a discovery (if NP exist)
- The LHC experiments have established an impressive variety of very powerful direct searches for many different final states!
 - Based on these results we need to establish the “big picture” in order to understand find out if/where our search strategy might have weak spots or even holes!
 - This requires appropriate interpretations of the searches and a MEANINGFUL comparison with other experiments – important example DM searches!
- The high energy running of the LHC starting 2015 will be our next very (as in very) real chance for discovery!


The story continues ... stay tuned!


BACKUP











ATLAS & CMS public results

All results presented in this talk (and many more)
can be accessed via the public page of the
ATLAS and CMS experiments:

ATLAS SUSY: <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>

CMS SUSY: <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>


ATLAS Summary

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: EPS 2013

ATLAS Preliminary


$$\int \mathcal{L} dt = (4.4 - 22.9) \text{ fb}^{-1} \quad \sqrt{s} = 7, 8 \text{ TeV}$$



*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.

CMS Summary

Summary of CMS SUSY Results* in SMS framework EPSHEP 2013



Summary

Slide from 2007
EPS Plenary talk in Manchester
LHC Detectors
Commissioning & Physics
pre-accident & 14 TeV assumption



- LHC&Experiments are on track for first collisions in 2008
 - Challenge: commissioning of machine and detectors of unprecedented complexity, technology and performance
- The LHC will discover (or exclude) the Higgs by ~2010
 - Electro Weak Symmetry Breaking
 - Large phase space can already be excluded with only $\sim 1\text{fb}^{-1}$
- The LHC will discover low energy SUSY (if it exists)
 - Could be easy; could also take more time and ingenuity before we can claim a discovery
 - First signals might emerge already in the first data
 - 1-3 TeV can be covered already with $< 10\text{fb}^{-1}$
- The LHC will cover a new physics scale of 1-3 TeV
 - Many new physics models; Black hole, Extra Dimensions, Little Higgs, Split Susy, New Bosons, Technicolour, etc ...

In other words; the next five years will be an exciting time for particle physics ...

(Best) mass limits in a nutshell (RP conserving)

SUSY & DM Searches, @ LHC O. Buchmüller

Direct squark	$\tilde{q} \rightarrow q\chi_1^0$	$\tilde{u}_L \rightarrow q\chi_1^0$	$\tilde{b} \rightarrow b\chi_1^0$	$\tilde{t} \rightarrow t\chi_1^0$
Best limit: [GeV]	~850	~500	~650	~650
No limit for M_{LSP} [GeV]	~300	~120	~270	~260

coloured sparticle production


Direct squark	$\tilde{g} \rightarrow q\bar{q}\chi_1^0$	$\tilde{g} \rightarrow b\bar{b}\chi_1^0$	$\tilde{g} \rightarrow t\bar{t}\chi_1^0$
Best limit: [GeV]	~1200	~1200	~1400
No limit for M_{LSP} [GeV]	~480	~650	~700

Stop $M_{\text{stop}} - M_{\text{LSP}} < M_{\text{top}}$	$\tilde{t} \rightarrow c\chi_1^0$	$\tilde{t} \rightarrow Wb\chi_1^0$
Best limit: [GeV]	~240	~320
No limit for M_{LSP} [GeV]	~210	~190







EWK sparticle production

Direct slepton	$\tilde{l}_L \rightarrow l^\pm\chi_1^0$	$\tilde{l}_R \rightarrow l^\pm\chi_1^0$
Best limit: [GeV]	~300	~240
No limit for M_{LSP} [GeV]	~150	~90





$\chi_1^\pm\chi_2^0$	light $\tilde{l}$	heavy $\tilde{l}$
Best limit: [GeV]	~750	~300
No limit for M_{LSP} [GeV]	~350	~60



Many Different Kinematic Variables



Multijets & missing energy search



Traditional inclusive Jets + E_T^{mis} search, which uses simple kinematic variables to categorize the events.

Main backgrounds QCD, W/Z+jet and $t\bar{t}$ are estimated using data-driven techniques.




CMS-SUS-PAS-13-012

See parallel talk for details:


C. Autermann



2010: Entering New Territory at the LHC!




Phys. Lett. B698:196-218




PLB 701 (2011) 186

Simplified Model Spectra (SMS)


Searches, @ LHC O. Buchmüller



$$g\tilde{g} \rightarrow q\bar{q}\tilde{\chi}^0 q\bar{q}\tilde{\chi}^0$$





$$q\bar{q} \rightarrow q\tilde{\chi}^0 q\bar{q}\tilde{\chi}^0$$




$$\tilde{q} \rightarrow q\tilde{\chi}_1^0, \tilde{g} \rightarrow q q\tilde{\chi}_1^0$$


CMS Preliminary, 19.5 fb⁻¹, $\sqrt{s} = 8$ TeV



$$m_G^{\text{max}} \approx 1.7 \text{ TeV}$$





Today: many more SMS and many more searches





Example: $\tilde{g}\tilde{g} \rightarrow t\bar{t}t\bar{t}\chi_1^0\chi_1^0$

Several searches are interpreted in this particular SMS!



(Minimal) Natural SUSY Spectrum

hmüller




Use the argument of “naturalness” (i.e. fine-tuning) to motivate light **3rd generation squarks** (especially stop) and a rather light **gluino**!

More in Gian Guidice talk!

What do we call a “SUSY search”?

*The definition is purely derived from the experimental signature.
Therefore, a “SUSY search signature” is characterized by*

Lots of missing energy, many jets, and possibly leptons in the final state



Missing Energy:

- from LSP

Multi-Jet:

- from cascade decay (gaugino)

Multi-Leptons:


- from decay of charginos/neutralinos

RP-Conserving SUSY is a very prominent example predicting this famous signature but ...

What is its experimental signature?

SUSY & DM Searches, @ LHC O. Buchmüller

... by no means is it the only New Physics model predicting this experimental pattern. Many other NP models predict this genuine signature



Missing Energy:

- N_{wimp} - end of the cascade

Multi-Jet:

- from decay of the N s (possibly via heavy SM particles like top, W/Z)


Multi-Leptons:

- from decay of the N 's


Model examples are Extra dimensions, Little Higgs, Technicolour, etc
but a more generic definition for this signature is as follows.

Rediscovery of the SM at a new energy frontier

SUSY & DM Searches, @ LHC O. Buchmüller



Combination vs individual search





Combination of
searches stable

Individual searches
exhibit large
variations

**Combinations is stable vs.
complexity while individual
searches are NOT!**

Natural SUSY: universal limits

If the gluino mass OR 3G mass lies in the red band, the point is excluded.
If the gluino mass AND 3G mass lie in the yellow band the point may or may not be excluded. Otherwise the point is not excluded.



Combining with the latest published 8 TeV results:

Outlook: 8 TeV vs 14 TeV

Use 30/fb for 2011/2012 for comparison

Higgs:

pp $\rightarrow$ H, $H \rightarrow WW, ZZ$ and $\gamma\gamma$
mainly gg: factor ~ 2

SUSY – 3rd Generation:

Mass scale ~ 500 GeV
qq and gg: factor ~ 3 to 6

SUSY – Squarks/Gluino:

Mass scale ~ 1.5 TeV
qq,gg,qq: factor ~ 40 to 80

Z' :

Mass scale ~ 5 TeV
qq: factor ~ 1000

Higgs:

15/fb@14 TeV to match 2011/2012
mainly gg: factor ~ 2

SUSY – 3rd Generation:

5/fb to 10/fb@14 TeV to match 2011/2012
qq and gg: factor ~ 3 to 6

SUSY – Squarks/Gluino:

0.4/fb to 0.8/fb@14 TeV to match 2011/2012
qq,gg,qq: factor ~ 40 to 80

Z' :

0(1/pb) @14 TeV to match 2011/2012
qq: factor ~ 1000

RP-violating searches/interpretation




Searches, @ LHC O. Buchmüller

Generic same-sign di-lepton
search with different signal regions

SR	Expected	Observed
RPV0	38 $\pm$ 14	35
RPV2	5.3 $\pm$ 2.1	5
SStop1	160 $\pm$ 59	152
SStop1++	90 $\pm$ 32	92
SStop2	40 $\pm$ 13	52
SStop2++	22 $\pm$ 8	25


$\tilde{g} \rightarrow 3 \text{ jets}$

Take all triplets,
QCD: $M_{3j} \sim \Sigma p_T^j$;
SUSY: $M_{3j} \sim M_g$
 $M_{3j} < \Sigma p_T^j - 160 \text{ GeV}$



RP violation searches: Summary



SUSY & DM Searches, @ LHC O. Buchmüller



ATLAS
similar

Like RP conserving searches, these searches are also probing the
1 TeV scale and even beyond!

Long-lived particle (SUSY) searches



ATLAS-CONF
-2013-069


nearly mass-degenerate

$$\chi_1^\pm \chi_1^0$$

search based on
disappearing-track
signature

CMS similar
See parallel talk from
L. Quertenmont

Long-lived particles	Direct $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^\pm$	Disapp. trk	1 jet	Yes	20.3
	Stable, stopped $\tilde{g}$ R-hadron	0	1-5 jets	Yes	22.9
	GMSB, stable $\tilde{\tau}$, $\tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\tilde{e}, \tilde{\mu}) + \tau(e, \mu)$	1-2 μ	0	-	15.9
	GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	0	Yes	4.7
	$\tilde{\chi}_1^0 \rightarrow qq\mu$ (RPV)	1 μ	0	Yes	4.4







$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) = 160 \text{ MeV}$, $\tau(\tilde{\chi}_1^\pm) = 0.2 \text{ ns}$
 $m(\tilde{\chi}_1^0) = 100 \text{ GeV}$, $10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$
 $10 < \tan\beta < 50$
 $0.4 < \tau(\tilde{\chi}_1^0) < 2 \text{ ns}$
 $1 \text{ mm} < c\tau < 1 \text{ m}$, $\tilde{g}$ decoupled

ATLAS-CONF-2013-069
 ATLAS-CONF-2013-057
 ATLAS-CONF-2013-058
 1304.6310
 1210.7451

About to probe
the 1 TeV scale


1 TeV

Dedicated searches for direct stop-pair production







Scalar stop analysis with two leptons in the final state using a MVA technique.




Dedicated searches for direct stop-pair production



LEP:
LEPSUSY
WG/04-01.1



CDF:
1203.4171

D0:
0803.2263



SMS limits: A word of caution!

SUSY & DM Searches, @ LHC O. Buchmüller



NS0	NS1	NS2	NS3	NS4
$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$
$\tilde{t}_1, \tilde{t}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$
		$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$
		$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm, \tilde{\ell}_{L,R}$
$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$

Used inclusive searches from 2011:

0-Lepton CMS-SUS-11-022



1-Lepton CMS-SUS-12-010

2-Lepton SS CMS-SUS-11-010

2-Lepton OS CMS-SUS-11-011

Combining Searches = less model dependence

SUSY & DM Searches, @ LHC O. Buchmüller



NS0	NS1	NS2	NS3	NS4
$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$
$\tilde{t}_1, \tilde{t}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$
		$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$
		$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm, \tilde{\ell}_{L,R}$
$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$

Used inclusive searches from 2011:

0-Lepton CMS-SUS-11-022


1-Lepton CMS-SUS-12-010

2-Lepton SS CMS-SUS-11-010

2-Lepton OS CMS-SUS-11-011

Combining Searches = less model dependence

SUSY & DM Searches, @ LHC O. Buchmüller



NS0				
$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$	$\tilde{g}$
$\tilde{t}_1, \tilde{t}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$	$\tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$
		$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$	$\tilde{\chi}_0^2$
		$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm$	$\tilde{\chi}^\pm, \tilde{\ell}_{L,R}$
$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$	$\tilde{\chi}_0^1$


2011:

0-Lepton CMS-SUS-11-022
1-Lepton CMS-SUS-12-010
2-Lepton SS CMS-SUS-11-010
2-Lepton OS CMS-SUS-11-011


Validity of Effective Field Theory Limits

Recent work from OB, M.Dolan,C.McCabe: arXiv:1308.6799

➤ Compare Effective Field Theory (EFT) with Full Theory (FT)



Use vector and axial-vector mediators (e.g. Z') as example - scalar are similar in conclusion!



Three Regions as function of mediator mass:

Region I: Heavy m_{med}

➤ EFT is valid!


Region II: Medium m_{med} – Resonant enhancement

➤ EFT limits are too conservative!

Region III: Low m_{med}

➤ EFT limits are too aggressive!

What those this imply on model-dependences of EFT limits?



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Must require $m_{\text{med}} < \Gamma_{\text{med}}$
2. Region in which EFT is valid (20%)
3. Require compatibility with relic density
4. Require theory to be perturbative ($< 4\pi$)

When we also require that the region/theory must be perturbative:

$$\sqrt{g_q g_\chi} < 4\pi$$


only a very small region is left!

This together with the observation that all DM theories for which the EFT is valid must have $m_{\text{med}} < \Gamma_{\text{med}}$ leads to the conclusion the the EFT only applies to a very (as in VERY) small class of DM models.

EFT limits of monojet searches are therefore highly model-dependended!

Why is SUSY so attractive?


1. Quadratically divergent quantum corrections to the Higgs boson mass are avoided



(Hierarchy or naturalness problem)

2. Unification of coupling constants of the three interactions seems possible


$$m_{\text{SUSY}} \sim 1 \text{ TeV}$$



3. SUSY provides a candidate for dark matter,



**The lightest
SUSY particle
(LSP)**



4. A SUSY extension is a small perturbation, consistent with the electroweak precision data


A “typical” SUSY Spectrum

Use the famous SPS1a benchmark point for illustration
 $[m_0=100, m_{1/2}=250, \tan\beta=10, A_0=-100, \mu>0]$

CMSSM

$m_0, m_{1/2}, \tan\beta, A_0, \text{sign}(\mu)$

SUSY & DM Searches, @ LHC O. Buchmüller



Advantage:


- Only four free parameters (when $\text{sign}(\mu)$ fixed)
- One of the most studied incarnations of the MSSM

Disadvantage:

- Not generally representative of SUSY (e.g. fixed mass relation between M_{gluon} and M_{LSP})

Stop decay to charm and neutralino


SUSY & DM Searches, @ LHC O. Buchmüller




ATLAS-CONF-2013-068:

Two different selections:

- Monojet-like selection to cover region close to 'diagonal'
- MVA based c-tag selection for remaining region




c-Tag:
95% excl.
visible XS:
 $\epsilon\sigma = 0.7 \text{ fb}$
signal obs:
13 events
signal exp:
 14^{+5}_{-4} events
 CL_B :
0.45



Monojet
95% excl.
visible XS:
 $E\sigma = 136 \text{ fb}$
signal obs:
2770 events
signal exp:
 2060^{+780}_{-560}
 CL_B :
0.86


Rediscovery of the SM at a new energy frontier

SUSY & DM Searches, @ LHC O. Buchmüller




Dark Matter from invisible Higgs searches


SUSY & DM Searches, @ LHC O. Buchmüller




Example CMS



CMS Preliminary



$\text{BR}(H \rightarrow \text{invisible}) < 68\% \text{ @ } 95\% \text{CL}$



$\text{BR}(H \rightarrow \text{invisible}) < 75\% \text{ @ } 95\% \text{CL}$

Dark Matter from invisible Higgs searches



SUSY & DM Searches, @ LHC O. Buchmüller

Status 2012 CMS only:

VBF: $BR_{H \rightarrow \text{invisible}} < 68\% @ 95\%CL$


VH: $BR_{H \rightarrow \text{invisible}} < 75\% @ 95\%CL$

Naïve combination: $\sim 50\% @ 95\% CL$



Assuming the experiments are able to maintain trigger and analysis acceptances, the LHC will provide a VERY powerful comparison of indirect & direct measurement of $\Gamma_{H \rightarrow \text{invisible}}$. In the (near) future this might provide a stringent constraint for $M_{DM} < M_H/2$


What those this imply on model-dependences of EFT limits?



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Region in which EFT is valid


For this we calculate the minimum coupling

$$\sqrt{g_q g_\chi} = m_{\text{med}} / \Lambda_{\text{CMS}}$$

that the simplified model must have for the EFT limits to apply. This is defined by region I (i.e. better than 20% agreement of FT and EFT).

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits




Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density

When exclude the region in which relic abundance is larger than the observed value of $\Omega_{\text{xx}} h^2 = 0.119$ only mediator masses above a few hundred GeV fulfill this.

* Coupling chose such that CMS EFT limit on Λ applies to FT

Model-dependences of EFT limits



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Must require $m_{\text{med}} < \Gamma_{\text{med}}$
2. Region in which EFT is valid (20%)
3. Require compatibility with relic density
4. Require theory to be perturbative ($< 4\pi$)


When we also require that the region/theory must be perturbative:

$$\sqrt{g_q g_\chi} < 4\pi$$

only a very small region is left!

EFT limits of monojet searches only apply to a very (as in VERY) small class of DM models!

Model-dependences of EFT limits



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density
3. Require theory to be perturbative ($< 4\pi$)
4. $m_{\text{med}} < \Gamma_{\text{med}}$ ALWAYS!

We also find that for all DM models the EFT is valid the mass of the mediator must be smaller than its width!


In the remaining part of the plot:

$$\sqrt{g_q g_\chi} > 2$$

a particle-like interpretation of the mediator is doubtful because of $m_{\text{med}} < \Gamma_{\text{med}}$!

See discussion about equation 3.5 in arXiv:1308.6799 for further details.

What those this imply on model-dependences of EFT limits?



Look at EFT validity in $m_{\text{DM}} - \text{coupling}^*$ plane!

1. Region in which EFT is valid (20%)
2. Require compatibility with relic density
3. Require theory to be perturbative ($< 4\pi$)
4. $m_{\text{med}} < \Gamma_{\text{med}}$ ALWAYS!


The observation that all DM theories for which the EFT is valid must have $m_{\text{med}} < \Gamma_{\text{med}}$ and the small class to models it applies in any case leads to the conclusion the EFT only applies to a very small class of DM models.

EFT limits of monojet searches are therefore highly model-dependent!

Alternative Interpretation Ansatz: Simplified models

Recent work from OB, M.Dolan, C.McCabe: arXiv:1308.6799



- Compare Effective Field Theory (EFT) with Full Theory (FT)



After three years of operation at the LHC the landscape for interpretation of searches has changed dramatically – new superior & modern approaches have replaced in many areas longstanding traditional ones (e.g. SUSY searches)



The proposal

SUSY & DM Searches, @ LHC O. Buchmüller





The proposal

SUSY & DM Searches, @ LHC O. Buchmüller





The proposal

SUSY & DM Searches, @ LHC O. Buchmüller



The proposal


SUSY & DM Searches, @ LHC O. Buchmüller



Very similar to limits
in SUSY simplified
models –
 m_{med}, m_{DM}, Λ
and possibly
some variation
of Γ_{med}
will cover the full
problem!

$pp \rightarrow \tilde{q} \tilde{q}, \tilde{q} \rightarrow q \tilde{\chi}_1^0; m(\tilde{g}) \gg m(\tilde{q})$
 --- Expected Limit $\pm 1 \sigma$ exp.
 $\sigma^{\text{NLO+NLL}} \pm 1 \sigma$ theory
 — $\tilde{q}_L + \tilde{q}_R, \tilde{u} + \tilde{d} + \tilde{s} + \tilde{c}$
 - - - $\tilde{u}_L$ only
 CMS, 11.7 fb^{-1}
 $\sqrt{s} = 8 \text{ TeV}$

If CMS is interested, we would be available to help with the implementation of simplified models for the monojet search!




124


Beyond EFT limits: Simplified models

Buchmüller

Working out the complementarity between direct DM detection experiments and collider based DM searches!



EFT limits give the impression that monojet searches outperform direct detection BUT EFT only applies a VERY small class of DM models.




Simplified model limits give a much better Account of the REAL complementarity and thus seem superior for a comparison.


Beyond EFT limits: Simplified models

Buchmüller

Working out the complementarity between direct DM detection experiments and collider based DM searches!




EFT limits give the impression that monojet searches outperform direct detection BUT EFT only applies a VERY small class of DM models.




Simplified model limits give a much better Account of the REAL complementarity and thus seem superior for a comparison.

Monojet and Monophoton (plus E_T^{miss})



Monojet: hard jet + E_T^{miss}



Monophoton: hard photon + E_T^{miss}




Note different y scale




Monophoton: hard photon + E_T^{miss}

Monophoton: hard photon + E_T^{miss}



Note different y scale



Spin Dependent

—■— CMS (90% C)

— CDF

SIMPL E 201

COUPP 2011

- IceCube ($\gamma\gamma \rightarrow W^+W^-$)

■ ■ ■ ■ ■ Super-K I+II+III ($\gamma\gamma \rightarrow W^+W^-$)